

# FORAGER PRO 40 LIQUID

For Cattle on Pasture.

## GUARANTEED ANALYSIS

Crude Protein*	not less than	40.0	%
* (This includes not more than 34% equivalent crude protein from non-protein nitrogen.)			
Crude Fiber	not more than	1.0	%
Calcium	not less than	0.1	%
Calcium	not more than	0.5	%
Phosphorus	not less than	1.0	%
Salt	not less than	2.0	%
Salt	not more than	2.5	%
Magnesium	not less than	0.3	%
Potassium	not less than	1.0	%
Copper	not less than	230	ppm
Iodine	not less than	20	ppm
Selenium	not less than	2.0	ppm
Zinc	not less than	760	ppm
Vitamin A	not less than	30,000	IU/lb
Vitamin D	not less than	3,000	IU/lb
Vitamin E	not less than	50	IU/lb
Total Sugars	not less than	20	%
Moisture	not more than	43	%

## INGREDIENTS

Condensed Glutamic Acid Fermentation Solubles, Molasses Products, Corn Syrup, Condensed Fermented Corn Extractives, Condensed Whey, Urea, Ammonium Polyphosphate, Water, Magnesium Oxide, Vitamin E Supplement, Sodium Selenite, Vitamin A Supplement, Vitamin D3 Supplement, Calcium Carbonate, Zinc Sulfate, Manganese Sulfate, Basic Copper Chloride, Ethylenediamine Dihydriodide, Cobalt Carbonate.

## FEEDING DIRECTIONS

**FOR FREE CHOICE FEEDING:** Cattle should consume 1 to 1.5 pounds per head per day depending on their nutritional needs and the quality of other feeds. Offer this product continuously in a lick wheel feeder. Moving the feeder closer to or farther from the water source or the area where cattle normally congregate will help control consumption. Do not feed to hungry or starving cattle. Forages fed with this product should generally be of good quality and offered free choice so that they will adequately supply the remainder of the animal's protein and energy sources.

**FOR CONTROLLED FEEDING:** Feed to cattle at the rate of 1 to 1.5 pounds per head per day. Thoroughly mix with grain and/or forage prior to feeding.

Free choice salt and fresh, clean water should be available at all times. The need for supplemental free choice mineral feeding will depend upon the kind and amount of ration ingredients being utilized. Ask your dealer and/or feed consultant for assistance in planning a feeding program to meet your specific needs. **RECIRCULATE OR AGITATE PRIOR TO USE.**

**Recirculation Systems:** Recirculate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top.

**Agitation Systems(mechanical, air, etc.):** Agitate immediately prior to use for not less than 10 minutes, creating turbulence at the bottom of the tank that is visible at the top of the tank.

Recirculate or agitate daily even when the supplement is not in use.

**CAUTION:** For ruminant use only. Do not feed to sheep.

**2350.**

Manufactured By:  
Midwest Ag Supplements, LLC  
Marshall, MN 56258

**Bulk Only Liquid**

This feed was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.